

Bijenhouden 2021 – nr. 1 februari

Biodiversiteit (6): Het belang van natuur in landbouwgebieden – Henk van der Scheer en Tjeerd Blacquièrre

Bastin, J.-F., Finegold, Y., Garcia, C., Mollicone, O., Rezende, M., Routh, O., Zohner, C.M. en Crowther, T.W., 2019. The global tree restoration potential. *Science* 365(6448):76-79.

Bouma, E., 2009. Klimaatverandering en plantgezondheid. *Gewasbescherming* 40(5):229-237.

Bruinsma, W., 2020. Natuurgebieden nabij landbouwgebieden zijn goed voor de bestuiving. *Bijenhouden* 14(6):32-33.

Challinor, A.J., Ewert, F., Arnold, S., Simelton, E. en Fraser, E., 2009. Crops and climate change: progress, trends, and challenges in simulating impacts and informing adaptation. *Journal of Experimental Botany* 60(10):2775-2789.

Garibaldi, L.A., Steffan-Dewenter, I., Kremen, C., Morales, J.M., Bommarco, R., Cunningham, S.A., Carvalheiro, L.G., Chacoff, N.P., Dudenhöffer, J.H., Greenleaf, S.S., Holzschuh, A., Isaacs, R., Krewenka, K., Mandelik, Y., Mayfield, M.M., Morandin, L.A., Potts, S.G., Ricketts, T.H., Szentgyörgyi, M., Viana, B.F., Westphal, C., Winfree, R. en Klein, A.-M., 2011. Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters* 14:1062-1072.

Gathmann, A en Tschardtke, T., 2002. Foraging ranges of solitary bees. *Journal of Animal Ecology* 71:757-764.

Geiger, F., Bengtsson, J., Berendse, F., Weisser, W.W., Emmerson, M., Morales, M.B., Ceryngier, P., Liira, J., Tschardtke, T., Winqvist, C., Eggers, S., Bommarco, R., Pärt, T., Bretagnolle, V., Plantegenest, M., Clement, L.W., Dennis, C., Palmerd, C., Onate, J.J., Guerrero, I., Hawro, V., Aavik, T., Thies, C., Flohre, A., Hänke, S., Fischer, C., Goedhart, P.W. en Inchausti, P., 2010. Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic and Applied Ecology* 11:97-105.

Groot, G.A. de, Kats, R. van, Reemer, M., Sterren, D. van der, Biesmeijer, J.C. en Kleijn, D., 2015. De bijdrage van (wilde) bestuivers aan de opbrengst van appels en blauwe bessen. *Alterra-rapport 2636*, Pp. 76, ISSN 1566-7197.

Heemert, C. van, 2021. Tekort aan bestuivende insecten in de VS. *Bijenhouden* 15(1):23-25.

Hofmann, M.M., Fleischmann, A en Renner, S.S., 2020. Foraging distances in six species of solitary bees with body lengths of 6 to 15 mm, inferred from individual tagging, suggest 150 m-rule-of-thumb for flower strip distances. *Journal of Hymenoptera Research* 77:105-117.

Hoog, A van 't, 2003. Grazen in een droomlandschap. *Bionieuws* 13(21):11.

Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C. en Tschardtke, T., 2007. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B* 274:303-313.

Kohier, F., Verhulst, J., Klink, R. van en Kleijn, D., 2008. At what spatial scale do high-quality habitats enhance the diversity of forbs and pollinators in intensively farmed landscapes? *Journal of Applied Ecology* 45:753-762.

Morandin, L.A., Winston, M.L., Abbott, V.A. en Franklin, M.T., 2007. Can pastureland increase wild bee abundance in agriculturally intense areas? *Basic and Applied* (ook wel koolstofassimilatie of koolzuurassimilatie) 8:117-124.

Proesmans, W., Bonte, O., Smagghe, G., Meeus, I. en Verheyen, K., 2019. Importance of forest fragments as pollinator habitat varies with season and guild. *Basic and Applied Ecology* 34:95-107.

Sandom, C.J., Ejrnæs, R., Hansen, M.D.D. en Svenning, J.-C., 2014. High herbivore density associated with vegetation diversity in interglacial ecosystems. *Proceedings of the National Academy of Sciences of the United States of America* 111(11):4162-4167.

Velde, J. van de, 2019. 'Alles wat bijen nodig hebben, is een B&B'. *Fruitteelt* 109(22):10-11.

Venturina, E.M., Drummond, F.A., Hoshida, A.K., Dibble, A.C. en Stack, L.B., 2017. Pollination reservoirs for wild bee habitat enhancement in cropping systems: a review. *Agroecology and sustainable food systems* 41(2):101-142.

Zeilmaker, R., 2010. Extra CO<sub>2</sub> is voor planten een zegen. *Bionieuws* 20(2):8-9.