

Bijenhouden 2021 – nr. 4 augustus

Serie: Bestuiving (en kader van JPTHijse)

Voedingswaarde stuifmeel bepaalt door bestuiver en plant

Kees van Heemert

Cruden, R.,W. 2009. Pollen grain size, stigma depth, and style length : the relationships revisited. *Plant systematics and evolution* 278: 223-238.

Hao, K., Tjan, Z-X., Wang, Z-C, en Huang, S-Q. 2020. Pollengrain size associated with pollinator feeding strategy. <https://doi.org/10.1098/rspb.2020.1191>

Leonhardt, S., D. en Blüthgen, N. 2012. The same but different: pollen foraging in honeybee and bumblebee colonies. *Apidologie* 43: 449–464.

Roulston, T'ai., H. Cane, J.H., en Buchmann, S. L. 2000. What governs protein content of pollen: Pollinator preferences, Pollen-pistil interactions or phylogeny?

[https://doi.org/10.1890/0012-9615\(2000\)070\[0617:WGPCOP\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2000)070[0617:WGPCOP]2.0.CO;2).

Ruedenauer, F.A., Spaethe, J., Kooi, C. van der en Leonhardt, S. D. 2019. Pollinator or pedigree: which factors determine the evolution of pollen nutrients? *Oecologia* 191: 349-358.

Ruedenauer, F.A., Raubenheimer, D., Kessner-Beierlein, D., Noack, L., Spaethe, J. en Leonhardt, S., D. 2020. Best be(e) on low fat: linking nutrient perception, regulation and fitness. *Ecology Letters* 23: 545-554. 2016.

Scheer, H. van der. 2021. Giftige stoffen in stuifmeel. *Bijenhouden* nr.3. --- aanvullen

Vaudo, A., D., Patcha, H., M., Mortensen, D., A., Tookera, J., F. en Grozinger, C., M. 2016. Macronutrient ratios in pollen shape bumble bee (*Bombus impatiens*) foraging strategies and floral preferences. <https://doi.org/10.1073/pnas.1606101113>

Yeaman, R., L., Roulston, T., H., en Carr, D., E. 2014. Pollen quality for pollinators tracks pollen quality for plants in *Mimulus guttatus*. *Ecosphere* (7) Article 91:1-8.